

Table 1

Recommended Adult Immunization Schedule by Age Group, United States, 2025

Vaccine	19–26 years	27–49 years	50–64 years	≥65 years
COVID–19	1 or more doses of 2024–2025 vaccine (See Notes)			2 or more doses of 2024–2025 vaccine (See Notes)
Influenza inactivated (IIV3, ccIIV3) Influenza recombinant (RIV3)	1 dose annually			
Influenza inactivated (aIIV3; HD–IIV3) Influenza recombinant (RIV3)	Solid organ transplant (See Notes)			
Influenza live, attenuated (LAIV3)	1 dose annually			
Respiratory syncytial virus (RSV)	Seasonal administration during pregnancy (See Notes)		60 through 74 years (See Notes)	≥75 years
Tetanus, diphtheria, pertussis (Tdap or Td)	1 dose Tdap each pregnancy; 1 dose Td/Tdap for wound management (See Notes)			
Measles, mumps, rubella (MMR)	1 dose Tdap, then Td or Tdap booster every 10 years			
Measles, mumps, rubella (MMR)	1 or 2 doses depending on indication (if born in 1957 or later)			
Varicella (VAR)	2 doses (if born in 1980 or later)		2 doses	
Zoster recombinant (RZV)	2 doses for immunocompromising conditions (See Notes)			
Human papillomavirus (HPV)	2 or 3 doses depending on age at initial vaccination or condition	27 through 45 years		2 doses
Pneumococcal (PCV15, PCV20, PCV21, PPSV23)	See Notes			
Hepatitis A (HepA)	2, 3, or 4 doses depending on vaccine			
Hepatitis B (HepB)	2, 3, or 4 doses depending on vaccine or condition			
Meningococcal A, C, W, Y (MenACWY)	1 or 2 doses depending on indication (See Notes for booster recommendations)			
Meningococcal B (MenB)	19 through 23 years	2 or 3 doses depending on vaccine and indication (See Notes for booster recommendations)		
Haemophilus influenzae type b (Hib)	1 or 3 doses depending on indication			
Mpox	2 doses			
Inactivated poliovirus (IPV)	Complete 3-dose series if incompletely vaccinated. Self-report of previous doses acceptable (See Notes)			